arXiv:1207.7341v2 [cond-mat.str-el] 4 Aug 2012

Topological Hamiltonian as an Exact Tool for Topological Invariants

Zhong Wang[1] and Binghai Yan[2]
[1] Institute for Advanced Study, Tsinghua University, Beijing 100084, China
[2]Institute for Inorganic & Analytical Chemistry, Johannes Gutenberg University of Mainz, 55099 Mainz, Germany, EU

(Dated: August 7, 2012)

We propose the concept "topological Hamiltonian" for various topological insulators in interacting systems. Although a bad tool for calculating energy spectra, "topological Hamiltonian" is an exact tool in producing topological invariants. We also show that, unlike the topological Hamiltonian, the natural "effective Hamiltonian" for quasiparticles can produce wrong topological invariants, despite its ability to give us excellent energy spectra. We present a physical explanation for this counterintuitive phenomenon from the surface state picture, more precisely, we show that the topological Hamiltonian determines the existence or absence of gapless surface modes. This confirms the topological Hamiltonian instead of the mean field effective Hamiltonian as an exact tool for calculating topological invariants in interacting systems. This result has significant implications for analytical and numerical calculations of topological invariants in interacting systems, in particular, it shows that the conventional band structure calculations of topological invariant can fail.

PACS numbers: 73.43.-f,71.70.Ej,75.70.Tj

Introduction. Topological insulators are new states of matter whose bulk is gapped just like ordinary insulators, but the surface is robustly gapless[1–4]. In the noninteracting limit, the gapless surface states are protected by bulk topological invariants [5–14] which are defined in terms of the noninteracting Bloch band states. Such topological invariants are inapplicable to systems with electron-electron interaction, though many numerical calculations resort to various versions of "effective non-interacting Hamiltonian". It is worth emphasizing that there is no general justification for these extensions of noninteracting topological invariants to interacting cases, though they may have good chance to produce correct result if the interaction is weak. More recently, topological insulators with strong electron-electron interaction attracts much interest[15–31], it is therefore highly desirable to develop topological invariants for interacting insulators. Topological field theory description[11] was developed for such interacting topological insulators, however, a major question remained to be answered was to obtain explicit formulas for the quantized topological responses. It was proposed in Ref.[32] that a Z_2 topological invariants can be defined in terms of Green's function for interacting topological insulators in three spatial dimensions (3D). There are numerous recent works making wide use of Green's function[33–38]. More recently, exact topological invariants in terms of zero frequency Green's function were systematically proposed[39, 40] for various topological insulators including quantum Hall insulators and time reversal invariant topological insulators in 2D and 3D, which found interesting applications in various context[36, 41–45].

The applicability of the topological invariants expressed in terms of zero frequency Green's function is beyond the "renormalized bands" (or quasiparticle) picture. Generally the quasiparticle picture is not applicable to the calculation of topological invariants, because the concept of "quasiparticle" itself can break down in a strongly correlated system. In the present paper, let us focus on systems in which the quasiparticle picture works as a good approximation, then one may naively expect that an effective Hamiltonian describing quasiparticles will produce the correct topological invariants. A qusiparticle with an approximate energy ω and momentum $\mathbf{k}$ has a self energy $\Sigma(\omega, \mathbf{k})$, therefore, if we consider a "flat-band" insulator with all filled bands having the same renormalized energy (or quasiparticle spectrum) ω_0, then we may guess that $\Sigma(\omega_0, \mathbf{k})$ will be important for obtaining topological invariants. However, this expectation is incorrect, instead, only $\Sigma(0, \mathbf{k})$ appears in the exact topological invariants[40]. It is our purpose of the present paper to understand this counterintuitive result.

The central observation of the present paper is that the zero frequency Green's function tells us about the existence or absence of the gapless surface states, while the more conventional "effective Hamiltonian", though an accurate tool for bulk energy spectra, fails to do this. We will give an explicit example for this surprising phenomenon in the main text. This physical observation from surface states confirms the zero frequency Green's function approach as a crucial tool in topological invariants calculation for interacting systems, which will be of great use in searching for new materials for topological insulators in systems with strong electron-electron interaction. We coined the concept "topological Hamiltonian" (see the main text) for inverse Green's function at zero frequency , from which various topological invariants can be readily calculated.

The two main results of this paper can be summarized as follows. First, topological Hamiltonian give us bad bulk energy spectra (thus it is far from being a good effective Hamiltonian for the bulk), but it gives us exact topological invariants; second, and more notably, topological Hamiltonian determines the gapless surface states.

To conclude this introduction, we emphasize that all systems studied in the present paper are gapped in the bulk, though the Green's function method is also fruitful in gapless systems[46, 47], in which momentum space defects can be topologically protected.

Topological Hamiltonian and Effective Hamiltonian. Let us fix the notation first. The many body Hamiltonian is given as $H = H_0 + H_1$, where $H_0 = \sum_{\mathbf{k}} c^\dagger_{\mathbf{k}} h_0(\mathbf{k}) c_{\mathbf{k}} = \sum_{\mathbf{k},\alpha,\beta} c^\dagger_{\mathbf{k}\alpha}[h_0(\mathbf{k})]_{\alpha\beta} c_{\mathbf{k}\beta}$ is the free quadratic Hamiltonian(α,β run through all degree of freedom other than momenta, and the chemical potential has been absorbed into the definition of H_0) , and H_1 term describes the electron-electron interaction. Various (Matsubara, Retarded, etc) Green's functions are defined in the frequency-momentum space in the usual way[48]. For instance, the free fermion system has Matsubara Green's function $G(i\omega, \mathbf{k}) = 1/[i\omega - h_0(\mathbf{k})]$. For interacting systems, Green's functions are still determined by the many body Hamiltonian, though their explicit forms can be very complicated.

Recently, in Ref.[39, 40] various exact topological invariants expressed in terms of zero frequency Green's function were systematically defined and derived. The idea is outlined as follows. The zero frequency Green's function $G(i\omega = 0, \mathbf{k})$ is a Hermitian matrix, therefore, in the eigenvalue equation

$$G^{-1}(0, \mathbf{k})|\alpha(0, k)\rangle = \mu_\alpha(0, \mathbf{k})|\alpha(0, k)\rangle \tag{1}$$

all $\mu_\alpha(0, \mathbf{k})$ are real numbers. An eigenvector $|\alpha(0, k)\rangle$ with $\mu_\alpha(0, \mathbf{k}) > 0$ is called an "R-zero", and the space spanned by R-zeros are called "R-space". We can define Berry gauge field and Berry curvature in the R-space, and then define topological invariants just in the same way as the noninteracting cases. For example, for two-dimensional topological insulators with broken time reversal symmetry (namely the quantum Hall insulator), we have the "generalized Chern number"[40]

$$C_1 = \frac{1}{2\pi}\int d^2k \mathcal{F}_{xy} \tag{2}$$

where $\mathcal{F}_{ij} = \partial_i \mathcal{A}_j - \partial_j \mathcal{A}_i$, and $\mathcal{A}_i = -i\sum_\alpha \langle \mathbf{k}\alpha|\partial_{k_i}|\mathbf{k}\alpha\rangle$. Here $|\mathbf{k}\alpha\rangle$ runs through an orthonormal basis of the R-space. This simple generalized Chern number was shown[40] to be as powerful as the more complicated topological invariants defined using frequency integral[32, 46, 49]. Systematic generalizations of this zero frequency Green's function formula to topological insulators and superconductors in other symmetry classes can be found in Ref.[40, 42], and certain special case was studied in Ref.[50, 51] from different motivation.

Since $G^{-1}(0, \mathbf{k})$ plays an crucial role in our approach, let us define the "topological Hamiltonian" as

$$h_{\text{topo}}(\mathbf{k}) = -G^{-1}(0, \mathbf{k}) \tag{3}$$

which can also be written as

$$h_{\text{topo}}(\mathbf{k}) = h_0(\mathbf{k}) + \Sigma(0, \mathbf{k}) \tag{4}$$

following the Schwinger-Dyson equation. With this definition, the generalized Chern number given in Eq.(2) is equivalent to the Chern number of an noninteracting system with free Hamiltonian $h_0(\mathbf{k}) = h_{\text{topo}}(\mathbf{k})$. Other topological invariants such as Z_2 invariants can also be calculated from the topological Hamiltonian in similar way[40]. An naive understanding of h_{topo} is to think it as an effective Hamiltonian for quasiparticles, namely that h_{topo} is a formula for "renormalized energy band". As we will see, this understanding is incorrect. In fact, h_{topo} is far away from being an effective Hamiltonian, which is expected to properly describe the quasiparticle. One of the failures of h_{topo} to be an effective Hamiltonian is that its eigenvalues can be far away from the true energy spectra, as we will show below. The powerful aspect of topological Hamiltonian approach is its ability to produce topological invariants rather than energy spectra.

What is the natural effective Hamiltonian producing accurate energy spectra? Let us start from the self-consistent equations[52, 53] for quasiparticle spectra, which can be written in a simplified notation as

$$[h_0(\mathbf{k}) + \Sigma(\omega_\alpha, \mathbf{k})]|u_\alpha(\mathbf{k})\rangle = \omega_\alpha(\mathbf{k})|u_\alpha(\mathbf{k})\rangle \tag{5}$$

which can be more compactly written as

$$G^{-1}(\omega_\alpha, \mathbf{k})|u_\alpha(k)\rangle = 0 \tag{6}$$

Eq.(5) is a self-consistent equation with the self energy Σ depending on the spectra ω_α. It is worth to emphasize that it is $\Sigma(\omega_\alpha, \mathbf{k})$ instead of $\Sigma(0, \mathbf{k})$ that appears in Eq.(5). In fact, this simple observation was the initial motivation of the present paper. In a general interacting system, $\Sigma(0, \mathbf{k})$ can be far away from $\Sigma(\omega_\alpha, \mathbf{k})$, therefore the topological Hamiltonian approach using $\Sigma(0, \mathbf{k})$ is a bad tool for energy spectra, though it is an powerful tool for topological invariants. A natural question is whether we can define physically meaningful topological invariants in terms of $\Sigma(\omega_\alpha, \mathbf{k})$ or $G^{-1}(\omega_\alpha, \mathbf{k})$. As we will see in the following, this effective Hamiltonian approach turns out to be less fruitful than the topological Hamiltonian approach.

Let us proceed with this $\Sigma(\omega_\alpha, \mathbf{k})$ approach to topological invariants. It is natural to define a frequency dependent "effective Hamiltonian"

$$h_{\text{eff}}(\omega, \mathbf{k}) = h_0(\mathbf{k}) + \Sigma(\omega, \mathbf{k}) \tag{7}$$

Furthermore, it seems also natural to define Berry connection in terms of $|u_\alpha(\mathbf{k})\rangle$ instead of $|\alpha(0, \mathbf{k})\rangle$ given in Eq.(1), since $|u_\alpha(\mathbf{k})\rangle$ more faithfully represents the quasiparticle picture in the interacting system. Now there is a difficulty in doing so, namely that $\Sigma(\omega, \mathbf{k})$ is generally not a Hermitian matrix, but we will show that even if this difficulty can be circumvented, the topological invariant defined in this approach is still incorrect. Let us write $\Sigma(\omega, \mathbf{k}) = \Sigma_1(\omega, \mathbf{k}) + i\Sigma_2(\omega, \mathbf{k})$, with both Σ_1 and Σ_2 Hermitian. The assumption that quasiparticle picture is a good approximation for our system is equivalent to saying roughly that Σ_2 can be ignored compared to Σ_1, since a non-vanishing Σ_2 gives an imaginary part to the quasiparticles and implies finite lifetimes. Good quasiparticle approximation implies that the the inverse lifetime should be significantly smaller than the characteristic energy scale of the quasiparticle. Therefore, let us replace $\Sigma(\omega, k)$ by $\Sigma_1(\omega, k)$ in Eq.(5), namely

$$[h_0(\mathbf{k}) + \Sigma_1(\omega_\alpha, \mathbf{k})]|u_\alpha(\mathbf{k})\rangle = \omega_\alpha(\mathbf{k})|u_\alpha(\mathbf{k})\rangle \tag{8}$$

Here we encounter another difficulty, namely that $|u_\alpha(k)\rangle$ are generally not orthogonal. This is a relatively minor problem since we can obtain a orthonormal basis by the Gram-Schmidt process. For 2D quantum Hall systems, let us define the "quasiparticle Chern number" as

$$\tilde{C}_1 = \frac{1}{2\pi}\int d^2k\tilde{\mathcal{F}}_{xy} \tag{9}$$

where $\tilde{\mathcal{F}}_{ij} = \partial_i\tilde{\mathcal{A}}_j - \partial_j\tilde{\mathcal{A}}_i$, and $\tilde{\mathcal{A}}_i = -i\sum_\alpha\langle\bar{u}_\alpha(\mathbf{k})|\partial_{k_i}|\bar{u}_\alpha(\mathbf{k})\rangle$, in which $|\bar{u}_\alpha(\mathbf{k})\rangle$ is an orthonormal basis in the space spanned by those $|u_\alpha(\mathbf{k})\rangle$ with negative $\omega_\alpha(\mathbf{k})$. For the simple model to be studied in the next section, $|u_\alpha(\mathbf{k})\rangle$ themselves will be sufficient.

In the remaining parts of this paper, we will show by explicit example that Eq.(9) differs from the correct topological invariants given by Eq.(2), therefore it must be incorrect. We then explain the reason why Eq.(2) instead of Eq.(9) is the correct topological invariants, from the surface state picture.

Effective Hamiltonian can produce wrong topological invariant. In this section we will give an explicit example in which the effective Hamiltonian approach to topological invariants is incorrect, for instance, Eq.(9) can fail for quantum Hall system even if the quasiparticle picture is good. Let us begin with a two-dimensional (2D) model of quantum anomalous Hall effect[54, 55] with electron-electron interaction. Let us assume the free Hamiltonian part H_0 describes a two band model[55], with

$$h_0(k) = \sin k_x\sigma_x + \sin k_y\sigma_y + [m_0 + (2 - \cos k_x - \cos k_y)]\sigma_z \tag{10}$$

Instead of giving explicit form for electron-electron interaction terms H_1, in the following we will choose an ansatz for self energy $\Sigma(\omega, \mathbf{k})$, which is sufficient for our purpose.

Let us make a simple ansatz that $\Sigma(\omega, \mathbf{k}) \approx \Sigma_z(\omega, \mathbf{k})\sigma_z$ near $\mathbf{k} = 0$, in which case the main idea can be most readily appreciated. Near $\mathbf{k} = 0$ we have $h_{\text{eff}}(\omega, \mathbf{k}) \approx k_x\sigma_x + k_y\sigma_y + [m_0 + \Sigma_z(\omega, \mathbf{k} = 0)]\sigma_z$. Additionally, we assume that the interaction preserves particle-hole symmetry of the free fermion Hamiltonian H_0, implying that $\Sigma_z(\omega, 0) = \Sigma_z(-\omega, 0)$. From Eq.(8)(taking $\mathbf{k} = 0$) we can obtain the self-consistent equation

$$m_{\text{eff}} = m_0 + \Sigma_z(m_{\text{eff}}, \mathbf{k} = 0) \tag{11}$$

for the two eigenvalues $\pm m_{\text{eff}}$ of $h_{\text{eff}}(\omega, \mathbf{k} = 0)$. This effective mass will be crucial in calculating $\tilde{C}_1$. On the other hand, the topological mass is defined as

$$m_{\text{topo}} = m_0 + \Sigma_z(0, \mathbf{k} = 0) \tag{12}$$

Near $\mathbf{k} = 0$, we have $h_{\text{topo}}(\mathbf{k}) \approx k_x\sigma_x + k_y\sigma_y + m_{\text{topo}}\sigma_z$. Following Eq.(2), the correct topological invariants calculated form the topological Hamiltonian h_{topo} is given as

$$C_1 = \frac{1}{4\pi}\int d^2k\hat{\mathbf{n}}\cdot(\partial_{k_x}\hat{\mathbf{n}} \times \partial_{k_y}\hat{\mathbf{n}}) \tag{13}$$

where the integrating range is the Brillouin zone, $\mathbf{n} = (n_x, n_y, n_z)$ is extracted from $h_{\text{topo}} = \boldsymbol{\sigma}\cdot\mathbf{n}$, and $\hat{\mathbf{n}} = \mathbf{n}/|\mathbf{n}|$. As a concrete example, let us pick $m_0 = -0.2, \Sigma_z(0, \mathbf{k} = 0) = 0.1$, then $m_{\text{topo}} = -0.1$, from which it follows that $C_1 = -1$. If $\Sigma_z(0.1, \mathbf{k} = 0) = 0.3$ (as an example), then $m_{\text{eff}} = 0.1$ is self-consistently obtained from Eq.(11). It follows that $\tilde{C}_1$ determined from Eq.(9) is $\tilde{C}_1 = 0$. We knew that Eq.(2) produces the correct topological invariants C_1[40], therefore, Eq.(9) fails in this case because

$$\tilde{C}_1 \neq C_1 \tag{14}$$

Topological Hamiltonian determines topological surface states. In the previous sections we have shown that the mean field calculation with the natural effective Hamiltonian h_{eff} produces wrong topological invariants, on the other hand, the "topological Hamiltonian" $h_{\text{topo}}(\mathbf{k}) = -G^{-1}(0, \mathbf{k})$ produces the correct topological invariants. This may seem mysterious since h_{topo} is a bad tool for quasiparticle spectra. Why should we expect h_{topo} instead of h_{eff} to produce the correct topological invariants? The answer, as we will show, is the fact that the gapless surface states are determined by h_{topo}. The key idea is that the zero energy surface states, if exist, will "feel" the self energy $\Sigma(\omega = 0)$ because they have zero energy. In sharp contrast to h_{topo}, h_{eff} is not directly related to the existence or absence of surface modes, despite of the fact that they they tell us the energy spectra of the bulk . As a well known fact, the existence of topologically protected surface states is a holographical manifestation of the bulk topology, therefore, the connection between h_{topo} and surface modes firmly establishes the h_{topo} approach to topological invariants.

Now let us flesh out the above idea in the simple two-band model studied in the previous section. To this end, let us first assume that there are zero energy surface states, and then see what it implies. Consider a 2D system with coordinate (x, y)[see Fig.(1)]. What we shall show is that at the interface between two bulk states with different C_1 (calculated from topological Hamiltonian), there is gapless surface state. Let us suppose that the many-body Hamiltonian H is a function of x, namely that certain parameters (e.g. interaction strength) of H vary as functions of x, then the topological Hamiltonian h_{topo} is also a function of x. Assuming the form of h_{topo} given in the previous section, we now have $h_{\text{topo}}(x) \approx -i\sigma_x\partial_x + \sigma_yk_y + m_{\text{topo}}(x)\sigma_z$ since h_{topo} is now a function of x, and the translational symmetry along x-direction is broken. Because it will not affect our conclusion, we take $|m_{\text{topo}}(x)| << 1$ to simplify our analysis. Let us suppose

$$m_{\text{topo}}(x) \to M \ (x >> 0); \quad m_{\text{topo}}(x) \to -M \ (x << 0) \tag{15}$$

and m_{topo} smoothly interpolate between this two limits near $x = 0$ [see Fig.(1b)].



FIG. 1: The gapless mode on the edge. (a) The physical setting and the zero mode. (b) m_{topo} as a function of x.

Due to the translational symmetry in y direction, k_y is a good quantum number, thus we can study the $k_y = 0$ branch first. As we have emphasized, the topological Hamiltonian $h_{\mathrm{topo}}(\mathbf{k}) = -G^{-1}(0, \mathbf{k}) = h_0(\mathbf{k}) + \Sigma(0, \mathbf{k})$ is a bad effective Hamiltonian for the bulk because the true energy spectra are nonzero in the bulk; on the other hand, h_{topo} is the tailor-made effective Hamiltonian for the zero mode living at the interface. Because the zero mode has exact zero energy, the self energy is given exactly by $\Sigma(\omega = 0)$. In other word, locally the zero mode is described by an "effective Hamiltonian" $h_{\mathrm{topo}}(x)$. With this key observation, it is now most natural to study the zero mode solution as the eigenvalue problem of h_{topo}. Explicitly, we have

$$h_{\mathrm{topo}} \to -i\sigma_x\partial_x + \sigma_z m_{\mathrm{topo}}(x) \tag{16}$$

where we have set $k_y = 0$ as mentioned before. We have omitted terms in higher order of $\mathbf{k}$ since we are concerned with low energy modes. If $M > 0$ [M is defined in Eq.(15)], the solution of zero mode from Eq.(16) is readily obtained as

$$|\psi(x)\rangle_{\text{zero mode}} = \frac{1}{\sqrt{2}}\begin{pmatrix} 1 \\ i \end{pmatrix} \exp[-\int^x dx' m_{\mathrm{topo}}(x')] \tag{17}$$

The assumption of existence of this zero mode requires that $m_{\mathrm{topo}}(x) < 0$ when $x \to -\infty$ and $m_{\mathrm{topo}}(x) > 0$ when $x \to \infty$ (otherwise the wavefunction diverges)[56, 57], which is satisfied by this solution. If $M < 0$, then there is also a zero mode solution with $(1, -i)^T$ replacing the $(1, i)^T$ factor in Eq.(17). Note that the existence of zero mode is fully determined by the asymptotic behaviors of $h_{\mathrm{topo}}(x)$ as $|x| >> 0$, therefore it is determined by the bulk topology. From the above explicit calculation at the interface we have seen that when $m_{\mathrm{topo}}(x)$ changes sign at the interface near $x = 0$, there are exact zero modes. The above calculation looks in parallel with the non-interacting cases in which we start form h_0 instead of h_{topo}, however, the crucial difference is that in interacting cases we have a frequency-dependent self energy, and $\Sigma(\omega = 0)$ appears in a self-consistent manner.

From the bulk calculation given in the previous section, we have $C_1(M > 0) = 0$ and $C_1(M < 0) = -1$ (remember that $|M| << 1$), thus the topological invariants at the two sides of interface are different. To summarize the above calculations, we have shown in an explicit example that when the topological invariants calculated from h_{topo} are different at the two sides of an interface, there are gapless surface states. In experiment, one side of the interface is often the vacuum, which can be regarded as a topologically trivial insulator, therefore, "different topological invariants" at the two sides of interface means that the insulator has nontrivial topological invariant ($C_1 \neq 0$ in this quantum Hall example). The fact that topological Hamiltonian $h_{\mathrm{topo}}(\mathbf{k})$ determines the existence or absence of gapless surface mode is among the central result of this paper. It is equally notable that h_{eff} plays no role in the solution of zero energy surface mode. Shortly speaking, we have "h_{eff} for the bulk states and h_{topo} for the surface states", though both h_{eff} and h_{eff} are defined in the bulk. This is the deep reason why h_{topo} instead of h_{eff} determines the bulk topological invariants.

Conclusions. In conclusion, we have shown that the "topological Hamiltonian" h_{topo}, defined in terms of zero frequency Green's function, faithfully produces the topological invariants for interacting topological insulators, while the natural qausiparticle effective Hamiltonian h_{eff} fails to do this, though it gives much better energy spectra. The deep reason, as we have shown, is that h_{topo} directly tells us the information of gapless surface states, while h_{eff} fails in this respect. Although we only write down explicit formulas for 2D quantum Hall insulators, the same analysis can also be applied to topological insulators in other symmetry classes. It is also worth noting that the purpose of this paper is to propose and understand the topological Hamiltonian approach as an exact tool, while more realistic models in which this approach will find wide applications remain to be studied.

Let us conclude with the remark that in band structure calculations we are generally more concerned about quasiparticle spectra, namely that various versions of the effective Hamiltonian approach are preferred, but as we have shown in this paper, such calculation can fail in topological invariant calculations. The topological Hamiltonian approach, on the other hand, faithfully produces correct topological invariants, thus it is an crucial tool in searching for topological insulators with strong electron-electron interaction.

ZW thanks Zheng-Yu Weng for helpful discussions. ZW is especially grateful to Shou-Cheng Zhang for fruitful discussions and collaborations in previous work. ZW is supported by Tsinghua University (No. 553401001). BY is supported by the ERC Advanced Grant (291472).

---

[1] X. -L. Qi and S. -C. Zhang, Phys. Today **63**, No.1, 33 (2010).
[2] J. Moore, Nature **464**, 194 (2010).
[3] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
[4] X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
[5] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Phys. Rev. Lett. **49**, 405 (1982).
[6] C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 146802 (2005).
[7] J. E. Moore and L. Balents, Phys. Rev. B **75**, 121306 (2007).
[8] R. Roy, Phys. Rev. B **79**, 195322 (2009).
[9] L. Fu and C. L. Kane, Phys. Rev. B **74**, 195312 (2006).
[10] L. Fu, C. L. Kane, and E. J. Mele, Phys. Rev. Lett. **98**, 106803 (2007).
[11] X.-L. Qi, T. Hughes, and S.-C. Zhang, Phys. Rev. B **78**, 195424 (2008).
[12] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, Phys. Rev. B **78**, 195125 (2008).
[13] S. Ryu, A. Schnyder, A. Furusaki, and A. Ludwig, New J. Phys. **12**, 065010 (2010).
[14] Z. Wang, X.-L. Qi, and S.-C. Zhang, New J. Phys. **12**, 065007 (2010).
[15] S. Raghu, X.-L. Qi, C. Honerkamp, and S.-C. Zhang, Phys. Rev. Lett. **100**, 156401 (2008).
[16] A. Shitade, H. Katsura, J. Kune, X.-L. Qi, S.-C. Zhang, and N. Nagaosa, Phys. Rev. Lett. **102**, 256403 (2009).
[17] Y. Zhang, Y. Ran, and A. Vishwanath, Phys. Rev. B **79**, 245331 (2009).
[18] B. Seradjeh, J. E. Moore, and M. Franz, Phys. Rev. Lett. **103**, 066402 (2009).
[19] D. A. Pesin and L. Balents, Nat. Phys. **6**, 376 (2010).
[20] L. Fidkowski and A. Kitaev, Physical Review B **81**, 134509 (2010).
[21] R. Li, J. Wang, X.-L. Qi, and S.-C. Zhang, Nature Physics **6**, 284 (2010).
[22] M. Dzero, K. Sun, V. Galitski, and P. Coleman, Phys. Rev. Lett. **104**, 106408 (2010).
[23] S. Rachel and K. L. Hur, e-print arXiv p. 1003.2238 (2010).
[24] J. Wen, A. Rüegg, C.-C. J. Wang, and G. A. Fiete, Phys. Rev. B **82**, 075125 (2010).
[25] J. Maciejko, X.-L. Qi, A. Karch, and S.-C. Zhang, Phys. Rev. Lett. **105**, 246809 (2010), URL `http://link.aps.org/doi/10.1103/PhysRevLett.105.246809`.
[26] M. Levin and A. Stern, Phys. Rev. Lett. **103**, 196803 (2009), URL `http://link.aps.org/doi/10.1103/PhysRevLett.103.196803`.
[27] X. Zhang, H. Zhang, J. Wang, C. Felser, and S.-C. Zhang, Science **335**, 1464 (2012), arxiv: 1111.1267.
[28] M. Hohenadler, T. C. Lang, and F. F. Assaad, Phys. Rev. Lett. **106**, 100403 (2011).
[29] S.-L. Yu, X. C. Xie, and J.-X. Li, Phys. Rev. Lett. **107**, 010401 (2011), URL `http://link.aps.org/doi/10.1103/PhysRevLett.107.010401`.
[30] T. Neupert, L. Santos, S. Ryu, C. Chamon, and C. Mudry, Phys. Rev. Lett. **108**, 046806 (2012).
[31] T. Yoshida, S. Fujimoto, and N. Kawakami, Phys. Rev. B **85**, 125113 (2012), URL `http://link.aps.org/doi/10.1103/PhysRevB.85.125113`.
[32] Z. Wang, X.-L. Qi, and S.-C. Zhang, Phys. Rev. Lett. **105**, 256803 (2010).
[33] L. Wang, X. Dai, and X. C. Xie, Phys. Rev. B **84**, 205116 (2011).
[34] L. Wang, H. Jiang, X. Dai, and X. C. Xie, arxiv: cond-mat/1109.6292.
[35] V. Gurarie, Phys. Rev. B **83**, 085426 (2011).
[36] L. Wang, X. Dai, and X. C. Xie, arxiv: cond-mat/1203.1124.
[37] A. M. Essin and V. Gurarie, Phys. Rev. B **84**, 125132 (2011), URL `http://link.aps.org/doi/10.1103/PhysRevB.84.125132`.
[38] O. V. Yazyev, E. Kioupakis, J. E. Moore, and S. G. Louie, Phys. Rev. B **85**, 161101 (2012), URL `http://link.aps.org/doi/10.1103/PhysRevB.85.161101`.
[39] Z. Wang, X.-L. Qi, and S.-C. Zhang, Phys. Rev. B **85**, 165126 (2012), URL `http://link.aps.org/doi/10.1103/PhysRevB.85.165126`.
[40] Z. Wang and S.-C. Zhang, ArXiv e-prints (2012), cond-mat/1203.1028, PRX in press.
[41] A. Go, W. Witczak-Krempa, G. S. Jeon, K. Park, and Y. B. Kim, arxiv: cond-mat/1202.4460.
[42] Z. Wang and S.-C. Zhang, ArXiv e-prints (2012), 1204.3149.
[43] J. C. Budich, R. Thomale, G. Li, M. Laubach, and S.-C. Zhang, arxiv: cond-mat/1203.2928.
[44] J. C. Budich and B. Trauzettel, ArXiv e-prints (2012), 1207.1104.
[45] S. R. Manmana, A. M. Essin, R. M. Noack, and V. Gurarie, ArXiv e-prints (2012), 1205.5095.
[46] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford University Press,USA, 2003).
[47] P. Hořava, Phys. Rev. Lett. **95**, 016405 (2005), URL `http://link.aps.org/doi/10.1103/PhysRevLett.95.016405`.
[48] G. D. Mahan, *Many-Particle Physics* (Plenum Press, 2000).
[49] K. Ishikawa and T. Matsuyama, Z. Phys. C **33**, 41 (1986).
[50] G. E. Volovik, arxiv: cond-mat/1111.4627.
[51] G. E. Volovik, arxiv: cond-mat/0912.0502.
[52] M. S. Hybertsen and S. G. Louie, Phys. Rev. Lett. **55**, 1418 (1985), URL `http://link.aps.org/doi/10.1103/PhysRevLett.55.1418`.
[53] M. S. Hybertsen and S. G. Louie, Phys. Rev. B **34**, 5390 (1986), URL `http://link.aps.org/doi/10.1103/PhysRevB.34.5390`.
[54] F. D. M. Haldane, Phys. Rev. Lett. **61**, 2015 (1988).
[55] X.L. Qi, Y.S. Wu, and S.C. Zhang, Phys. Rev. B **74**, 085308 (2006).
[56] R. Jackiw and C. Rebbi, Phys. Rev. D **13**, 3398 (1976).
[57] J. Goldstone and F. Wilczek, Phys. Rev. Lett. **47**, 986 (1981).